

B-Surprise Putnam Problems

Each problem below is selected from the B session of the Putnam exam, from 2015 to 2019. These were not the easiest problems; rather they were 2nd or 3rd easiest, measured by the number of full-credit solutions received. See if you can make progress on any of these. These problems, solutions, and rankings are all from the Putnam Archive.

Problems:

2015 B–2. Given a list of the positive integers $1, 2, 3, 4, \dots$, take the first three numbers $1, 2, 3$ and their sum 6 and cross all four numbers off the list. Repeat with the three smallest remaining numbers $4, 5, 7$ and their sum 16 . Continue in this way, crossing off the three smallest remaining numbers and their sum, and consider the sequence of sums produced: $6, 16, 27, 36, \dots$. Prove or disprove that there is some number in the sequence whose base 10 representation ends with 2015 .

2016 B–2. Define a positive integer n to be *squarish* if either n is itself a perfect square or the distance from n to the nearest perfect square is a perfect square. For example, 2016 is squarish, because the nearest perfect square to 2016 is $45^2 = 2025$ and $2025 - 2016 = 9$ is a perfect square. (Of the positive integers between 1 and 10 , only 6 and 7 are not squarish.)

For a positive integer N , let $S(N)$ be the number of squarish integers between 1 and N , inclusive. Find positive constants α and β such that

$$\lim_{N \rightarrow \infty} \frac{S(N)}{N^\alpha} = \beta,$$

or show that no such constants exist.

2017 B–3. Suppose that $f(x) = \sum_{i=0}^{\infty} c_i x^i$ is a power series for which each coefficient c_i is 0 or 1 . Show that if $f(2/3) = 3/2$, then $f(1/2)$ must be irrational.

2018 B–2. Let n be a positive integer, and let $f_n(z) = n + (n-1)z + (n-2)z^2 + \dots + z^{n-1}$. Prove that f_n has no roots in the closed unit disk $\{z \in \mathbb{C} : |z| \leq 1\}$.

2019 B–5. Let F_m be the m th Fibonacci number, defined by $F_1 = F_2 = 1$ and $F_m = F_{m-1} + F_{m-2}$ for all $m \geq 3$. Let $p(x)$ be the polynomial of degree 1008 such that $p(2n+1) = F_{2n+1}$ for $n = 0, 1, 2, \dots, 1008$. Find integers j and k such that $p(2019) = F_j - F_k$.